

Universal acyclic resolutions for arbitrary coefficient groups

Michael Levin

Abstract

We prove that for every compactum X and every integer $n \geq 2$ there are a compactum Z of $\dim \leq n + 1$ and a surjective UV^{n-1} -map $r : Z \longrightarrow X$ such that for every abelian group G and every integer $k \geq 2$ such that $\dim_G X \leq k \leq n$ we have $\dim_G Z \leq k$ and r is G -acyclic.

Keywords: cohomological dimension, cell-like and acyclic resolutions

Math. Subj. Class.: 55M10, 54F45.

1 Introduction

This paper is devoted to proving the following theorem which was announced in [8].

Theorem 1.1 *Let X be a compactum. Then for every integer $n \geq 2$ there are a compactum Z of $\dim \leq n + 1$ and a surjective UV^{n-1} -map $r : Z \longrightarrow X$ having the property that for every abelian group G and every integer $k \geq 2$ such that $\dim_G X \leq k \leq n$ we have that $\dim_G Z \leq k$ and r is G -acyclic.*

The cohomological dimension $\dim_G X$ of X with respect to an abelian group G is the least number n such that $\check{H}^{n+1}(X, A; G) = 0$ for every closed subset A of X . A space is G -acyclic if its reduced Čech cohomology groups modulo G are trivial, a map is G -acyclic if every fiber is G -acyclic. By the Vietoris-Begle theorem a surjective G -acyclic map of compacta cannot raise the cohomological dimension $\dim_G$. A compactum X is approximately n -connected if any embedding of X into an ANR has the UV^n -property, i.e. for every neighborhood U of X there is a smaller neighborhood $X \subset V \subset U$ such that the inclusion $V \subset U$ induces the zero homomorphism of the homotopy groups in $\dim \leq n$. An approximately n -connected compactum has trivial reduced Čech cohomology groups in $\dim \leq n$ with respect to any group G . A map is called a UV^n -map if every fiber is approximately n -connected.

Theorem 1.1 generalizes the following results of [6, 7].

Theorem 1.2 ([6]) *Let G be an abelian group and let X be a compactum with $\dim_G X \leq n$, $n \geq 2$. Then there are a compactum Z with $\dim_G Z \leq n$ and $\dim Z \leq n + 1$ and a G -acyclic map $r : Z \rightarrow X$ from Z onto X .*

Theorem 1.3 ([7]) *Let X be a compactum with $\dim_{\mathbb{Z}} X \leq n \geq 2$. Then there exist a compactum Z with $\dim Z \leq n$ and a cell-like map $r : Z \rightarrow X$ from Z onto X such for every integer $k \geq 2$ and every group G such that $\dim_G X \leq k$ we have $\dim_G Z \leq k$.*

Theorem 1.2 obviously follows from Theorem 1.1. Theorem 1.3 can be derived from Theorem 1.1 as follows. Recall that a compactum is cell-like if any map from the compactum to a CW-complex is null-homotopic. A map is cell-like if its fibers are cell-like. Let X be of $\dim_{\mathbb{Z}} X < \infty$ and let $r : Z \rightarrow X$ satisfy the conclusions of Theorem 1.1 for $n = \dim_{\mathbb{Z}} X + 1$. Then $\dim_{\mathbb{Z}} Z \leq \dim_{\mathbb{Z}} X \leq n - 1$ and because Z is finite dimensional we have $\dim Z = \dim_{\mathbb{Z}} Z \leq n - 1$. Since r is UV^{n-1} and $\dim Z \leq n - 1$ we get that r is cell-like. Let for a group G , $\dim_G X \leq k \geq 2$. If $k \leq n$ then by Theorem 1.1, $\dim_G Z \leq k$ and if $k > n$ then $\dim_G Z \leq k$ since $\dim Z \leq n - 1$. Thus Theorem 1.1 implies Theorem 1.3.

It was observed in [7] that the restriction $k \geq 2$ in Theorem 1.3 cannot be omitted. Therefore Theorem 1.1 does not hold for $k = 1$.

Let us discuss possible generalizations of Theorem 1.1. One is tempted to reduce the dimension of Z to n . It is partially justified by

Theorem 1.4 ([8]) *Let X be a compactum. Then for every integer $n \geq 2$ there are a compactum Z of $\dim \leq n$ and a surjective UV^{n-1} -map $r : Z \rightarrow X$ having the property that for every finitely generated abelian group G and every integer $k \geq 2$ such that $\dim_G X \leq k \leq n$ we have that $\dim_G Z \leq k$ and r is G -acyclic.*

However Theorem 1.4 does not hold for arbitrary groups G . Indeed, one can show that a $\mathbb{Q}$ -acyclic UV^1 -map from a compactum of $\dim \leq 2$ must be $\mathbb{Z}$ -acyclic (even cell-like). Then a compactum X with $\dim_{\mathbb{Z}} X = 3$ and $\dim_{\mathbb{Q}} X = 2$ cannot be the image of a compactum of $\dim \leq 2$ under a $\mathbb{Q}$ -acyclic UV^1 -map.

The situation becomes more complicated if we drop in Theorem 1.1 the requirement that r is UV^{n-1} and consider

Problem 1.5 *Given a compactum X , an integer $n \geq 2$ and a collection of abelian groups $\mathcal{G}$ such that $\dim_G X \leq n$ for every $G \in \mathcal{G}$ do there exist a compactum Z of $\dim \leq n$ and a $\mathcal{G}$ -acyclic surjective map $r : Z \rightarrow X$ such that $\dim_G Z \leq \max\{\dim_G X, 2\}$ for every $G \in \mathcal{G}$? (The $\mathcal{G}$ -acyclicity means the G -acyclicity for every $G \in \mathcal{G}$.)*

In general the answer to Problem 1.5 is negative [5]. Indeed, let X be a compactum with $\dim_{\mathbb{Z}} = 3$, $\dim_{\mathbb{Q}} = 2$ and $\dim_{\mathbb{Z}_p} = 2$ for every prime p and let $G = \mathbb{Q} \oplus (\mathbb{Q}/\mathbb{Z})$. Clearly $\dim_G X = 2$ and the G -acyclicity implies both the $\mathbb{Q}$ and $(\mathbb{Q}/\mathbb{Z})$ -acyclicities. Then it follows from the Bockstein sequence generated by

$$0 \longrightarrow \mathbb{Z} \longrightarrow \mathbb{Q} \longrightarrow \mathbb{Q}/\mathbb{Z} \longrightarrow 0$$

that the G -acyclicity implies the $\mathbb{Z}$ -acyclicity and therefore there is no G -acyclic resolution for X from a compactum of $\dim \leq 2$.

The situation described in the example can be interpreted in terms of Bockstein Theory. Let $\mathcal{G}$ be a collection of abelian groups. Denote by $\sigma(\mathcal{G})$ the union of the Bockstein basises $\sigma(G)$ of all $G \in \mathcal{G}$. Based on the Bockstein inequalities define the closure $\overline{\sigma(\mathcal{G})}$ of $\sigma(\mathcal{G})$ as a collection of abelian groups containing $\sigma(\mathcal{G})$ and possibly some additional groups determined by:

$$\begin{aligned} \mathbb{Z}_p &\in \overline{\sigma(\mathcal{G})} \text{ if } \mathbb{Z}_{p^\infty} \in \overline{\sigma(\mathcal{G})}; \\ \mathbb{Z}_{p^\infty} &\in \overline{\sigma(\mathcal{G})} \text{ if } \mathbb{Z}_p \in \overline{\sigma(\mathcal{G})}; \\ \mathbb{Q} &\in \overline{\sigma(\mathcal{G})} \text{ if } \mathbb{Z}_{(p)} \in \overline{\sigma(\mathcal{G})}; \\ \mathbb{Z}_{(p)} &\in \overline{\sigma(\mathcal{G})} \text{ if } \mathbb{Q} \text{ and } \mathbb{Z}_{p^\infty} \in \overline{\sigma(\mathcal{G})}. \end{aligned}$$

One can show that for compact metric spaces the $\mathcal{G}$ -acyclicity implies the $\overline{\sigma(\mathcal{G})}$ -acyclicity. This motivates the following

Conjecture 1.6 *Problem 1.5 can be answered positively if $\dim_E X \leq n$ for every $E \in \overline{\sigma(\mathcal{G})}$.*

The key open case of this conjecture seems to be constructing a $\mathbb{Q}$ -acyclic resolution $r : Z \longrightarrow X$ for a compactum X with $\dim_{\mathbb{Q}} \leq n$, $n \geq 2$ from a compactum Z of $\dim \leq n$.

2 Preliminaries

All groups are assumed to be abelian and functions between groups are homomorphisms. $\mathcal{P}$ stands for the set of primes. For a non-empty subset $\mathcal{A}$ of $\mathcal{P}$ let $S(\mathcal{A}) = \{p_1^{n_1} p_2^{n_2} \dots p_k^{n_k} : p_i \in \mathcal{A}, n_i \geq 0\}$ be the set of positive integers with prime factors from $\mathcal{A}$ and for the empty set define $S(\emptyset) = \{1\}$. Let G be a group and $g \in G$. We say that g is $\mathcal{A}$ -torsion if there is $n \in S(\mathcal{A})$ such that $ng = 0$ and g is $\mathcal{A}$ -divisible if for every $n \in S(\mathcal{A})$ there is $h \in G$ such that $nh = g$. $\text{Tor}_{\mathcal{A}} G$ is the subgroup of the $\mathcal{A}$ -torsion elements of G . G is $\mathcal{A}$ -torsion if $G = \text{Tor}_{\mathcal{A}} G$, G is $\mathcal{A}$ -torsion free if $\text{Tor}_{\mathcal{A}} G = 0$ and G is $\mathcal{A}$ -divisible if every element of G is $\mathcal{A}$ -divisible.

G is $\mathcal{A}$ -local if G is $(\mathcal{P} \setminus \mathcal{A})$ -divisible and $(\mathcal{P} \setminus \mathcal{A})$ -torsion free. The $\mathcal{A}$ -localization of G is the homomorphism $G \longrightarrow G \otimes \mathbb{Z}_{(\mathcal{A})}$ defined by $g \longrightarrow g \otimes 1$ where $\mathbb{Z}_{(\mathcal{A})} = \{n/m : n \in \mathbb{Z}, m \in S(\mathcal{P} \setminus \mathcal{A})\}$. G is $\mathcal{A}$ -local if and only if the $\mathcal{A}$ -localization of G is an isomorphism. A simply connected CW-complex is $\mathcal{A}$ -local if its homotopy groups are $\mathcal{A}$ -local. A map between two simply connected CW-complexes is an $\mathcal{A}$ -localization if the induced homomorphisms of the homotopy and the (reduced

integral) homology groups are $\mathcal{A}$ -localizations.

The extensional dimension of a compactum X is said not to exceed a CW-complex K , written $\text{e-dim} X \leq K$, if for every closed subset A of X and every map $f : A \rightarrow K$ there is an extension of f over X . It is well-known that $\text{dim} X \leq n$ is equivalent to $\text{e-dim} X \leq \mathbb{S}^n$ and $\text{dim}_G X \leq n$ is equivalent to $\text{e-dim} X \leq K(G, n)$ where $K(G, n)$ is an Eilenberg-Mac Lane complex of type (G, n) .

A map between CW-complexes is said to be combinatorial if the preimage of every subcomplex of the range is a subcomplex of the domain. Let M be a simplicial complex and let $M^{[k]}$ be the k -skeleton of M (=the union of all simplexes of M of $\text{dim} \leq k$). By a resolution $EW(M, k)$ of M we mean a CW-complex $EW(M, k)$ and a combinatorial map $\omega : EW(M, k) \rightarrow M$ such that ω is 1-to-1 over $M^{[k]}$. Let $f : N \rightarrow K$ be a map of a subcomplex N of M into a CW-complex K . The resolution is said to be suitable for f if the map $f \circ \omega|_{\omega^{-1}(N)}$ extends to a map $f' : EW(M, k) \rightarrow K$. We call f' a resolving map for f . The resolution is said to be suitable for a compactum X if for every simplex Δ of M , $\text{e-dim} X \leq \omega^{-1}(\Delta)$. Note that if $\omega : EW(M, k) \rightarrow M$ is a resolution suitable for X then for every map $\phi : X \rightarrow M$ there is a map $\psi : X \rightarrow EW(M, k)$ such that for every simplex Δ of M , $(\omega \circ \psi)(\phi^{-1}(\Delta)) \subset \Delta$. We call ψ a combinatorial lifting of ϕ .

Let M be a finite simplicial complex and let $f : N \rightarrow K$ be a cellular map from a subcomplex N of M to a CW-complex K such that $M^{[k]} \subset N$. A standard way of constructing a resolution suitable for f is described in [7]. Such a resolution $\omega : EW(M, k) \rightarrow M$ is called the standard resolution of M for f and it has the following properties:

$EW(M, k)$ is $(k - 1)$ -connected if so are M and K ;

ω is a map onto and for every simplex Δ of M , $\omega^{-1}(\Delta)$ is either contractible or homotopy equivalent to K ;

for every subcomplex T of M , $\omega|_{\omega^{-1}(T)} : EW(T, k) = \omega^{-1}(T) \rightarrow T$ is the standard resolution of T for $f|_{N \cap T} : N \cap T \rightarrow K$.

Let G be a group, let $\alpha : L \rightarrow M$ be a surjective combinatorial map of a CW-complex L and a finite simplicial complex M and let n be a positive integer such that $\tilde{H}_i(\alpha^{-1}(\Delta); G) = 0$ for every $i < n$ and every simplex Δ of M . One can show by induction on the number of simplexes of M using the Mayer-Vietoris sequence and the Five Lemma that $\alpha_* : \tilde{H}_i(L; G) \rightarrow \tilde{H}_i(M; G)$ is an isomorphism for $i < n$. We will refer to this fact as the combinatorial Vietoris-Begle theorem.

Proposition 2.1 ([2]). *Let G be a group and $p \in \mathcal{P}$. The following conditions are equivalent:*

G is p -divisible;
 $\text{Ext}(\mathbb{Z}_{p^\infty}, G)$ is p -divisible;
 $\text{Ext}(\mathbb{Z}_{p^\infty}, G) = 0$.

Proposition 2.2 *Let G be a group, let $2 \leq k \leq n$ be integers and let $\mathcal{F} \subset \mathcal{P}$ and $p \in \mathcal{P} \setminus \mathcal{F}$. Let M be an $(n-1)$ -connected fine simplicial complex such that $H_n(M)$ is $\mathcal{F}$ -torsion and let $\omega : L = EW(M, k) \longrightarrow M$ be the standard resolution of M for a cellular map $f : N \longrightarrow K(G, k)$ from a subcomplex N of M containing $M^{[k]}$. Then L is $(k-1)$ -connected and for every $1 \leq i \leq n-1$*

- (i) $\pi_i(L)$ and $\pi_n(L)/\text{Tor}_{\mathcal{F}}\pi_n(L)$ are p -torsion if $G = \mathbb{Z}_p$;
- (ii) $\pi_i(L)$ and $\pi_n(L)/\text{Tor}_{\mathcal{F}}\pi_n(L)$ are p -torsion and $\pi_k(L)$ is p -divisible if $G = \mathbb{Z}_{p^\infty}$
- (iii) $\pi_i(L)$ and $\pi_n(L)/\text{Tor}_{\mathcal{F}}\pi_n(L)$ are q -divisible and $\pi_i(L)$ is q -torsion free for every $q \in \mathcal{P}$, $q \neq p$ if $G = \mathbb{Z}_{(p)}$;
- (iv) $\pi_i(L)$ and $\pi_n(L)/\text{Tor}_{\mathcal{F}}\pi_n(L)$ are q -divisible and $\pi_i(L)$ is q -torsion free for every $q \in \mathcal{P}$ if $G = \mathbb{Q}$.

Proof. Recall that ω is a combinatorial surjective map, for every simplex Δ of M , $\omega^{-1}(\Delta)$ is either contractible or homotopy equivalent to $K(G, k)$ and L is $(k-1)$ -connected because so are M and $K(G, k)$. Since M is $(n-1)$ -connected and $H_n(M)$ is $\mathcal{F}$ -torsion we have that $H_n(M; \mathbb{Q}) = 0$ and $H_n(M; \mathbb{Z}_q) = 0$, $H_n(M; \mathbb{Z}_{(q)}) = 0$ for $q \in \mathcal{P} \setminus \mathcal{F}$ and $H_n(M; \mathbb{Z}_{q^\infty}) = 0$ for every $q \in \mathcal{P}$.

(i) By the generalized Hurewicz theorem $\tilde{H}_*(K(\mathbb{Z}_p, k))$ is p -torsion. Then $\tilde{H}_*(K(\mathbb{Z}_p, k); \mathbb{Q}) = 0$. Hence by the combinatorial Vietoris-Begle theorem $\tilde{H}_i(L; \mathbb{Q}) = 0$ for $i \leq n$ and therefore $\tilde{H}_i(L)$ is torsion for $i \leq n$.

Let $q \in \mathcal{P}$ and $q \neq p$ and $i \leq n-1$.

Note $\tilde{H}_*(K(\mathbb{Z}_p, k); \mathbb{Z}_{(q)}) = 0$ and hence by the combinatorial Vietoris-Begle theorem $\tilde{H}_i(L; \mathbb{Z}_{(q)}) = 0$. Then $\tilde{H}_i(L) \otimes \mathbb{Z}_{(q)} = 0$. Thus $\tilde{H}_i(L)$ is torsion and q -torsion free and hence $\tilde{H}_i(L)$ is p -torsion.

Now let $q \in \mathcal{P} \setminus \mathcal{F}$ and $q \neq p$.

Recall $H_n(M; \mathbb{Z}_{(q)}) = 0$. Then using the previous argument we conclude that $H_n(L)$ is q -torsion free and hence $H_n(L)$ is $(\mathcal{F} \cup \{p\})$ -torsion.

By the generalized Hurewicz theorem $\pi_i(L)$ is p torsion for $i \leq n-1$ and $\pi_n(L)$ is $(\mathcal{F} \cup \{p\})$ -torsion. Thus $\pi_n(L)/\text{Tor}_{\mathcal{F}}\pi_n(L)$ is p -torsion and (i) follows.

(ii) The argument used in (i) applies to show that $\pi_i(L)$, $i \leq n-1$ and $\pi_n(L)/\text{Tor}_{\mathcal{F}}\pi_n(L)$ are p -torsion. Note that $\pi_k(L) = H_k(L)$. We will show that $H_k(L)$ is p -divisible and this will imply (ii). Observe that $H_k(K(\mathbb{Z}_{p^\infty}, k); \mathbb{Z}_p) = \mathbb{Z}_{p^\infty} \otimes \mathbb{Z}_p = 0$. Then since

$H_k(M; \mathbb{Z}_p) = 0$ the combinatorial Vietoris-Begle theorem implies that $H_k(L; \mathbb{Z}_p) = 0$. Thus $H_k(L) \otimes \mathbb{Z}_p = 0$ and therefore $H_k(L)$ is p -divisible.

(iii) Since $Z_{(p)}$ is p -local we have that $\tilde{H}_*(K(Z_{(p)}, k))$ is p -local and therefore $\tilde{H}_*(K(Z_{(p)}, k); \mathbb{Z}_q) = \tilde{H}_*(K(Z_{(p)}, k); \mathbb{Z}_{q^\infty}) = 0$ for every $q \in \mathcal{P}$, $q \neq p$.

Let $q \in \mathcal{P}$, $q \neq p$.

Recall that $\tilde{H}_i(M; \mathbb{Z}_{q^\infty}) = 0$ for $i \leq n$. Then by the combinatorial Vietoris-Begle theorem $\tilde{H}_i(L; \mathbb{Z}_{q^\infty}) = 0$ for $i \leq n$. Hence by virtue of the universal coefficient theorem $\tilde{H}_i(L) * \mathbb{Z}_{q^\infty} = 0$ and $\tilde{H}_i(L) \otimes \mathbb{Z}_{q^\infty} = 0$ for $i \leq n - 1$ and therefore $\tilde{H}_i(L)$ is q -torsion free and q -divisible for $i \leq n - 1$.

Let $q \in \mathcal{P}$, $q \neq p$ and $q \notin \mathcal{F}$.

Recall that $H_n(M; \mathbb{Z}_q) = 0$. By the combinatorial Vietoris-Begle theorem $H_n(L; \mathbb{Z}_q) = 0$. Hence $H_n(L) \otimes \mathbb{Z}_q = 0$ and therefore $H_n(L)$ is q -divisible.

Let $q \in \mathcal{P}$, $q \neq p$ and $q \in \mathcal{F}$.

Then $H_n(M; \mathbb{Z}_{q^\infty}) = 0$. By the combinatorial Vietoris-Begle theorem $H_n(L; \mathbb{Z}_{q^\infty}) = 0$. Hence $H_n(L) \otimes \mathbb{Z}_{q^\infty} = 0$ and therefore $H_n(L)/\text{Tor}_q H_n(L)$ is q -divisible.

Now using Completion and Localization Theories [1] we will pass to the homotopy groups of L .

Let $q \in \mathcal{P}$ and $q \neq p$.

Denote $\mathcal{A} = \mathcal{P} \setminus \{q\}$ and let $\alpha : L \longrightarrow L_\alpha$ be an $\mathcal{A}$ -localization of L . Recall that $\tilde{H}_i(L)$ is q -torsion free and q -divisible for $i \leq n - 1$. Then α induces an isomorphism of the groups $\tilde{H}_*(L)$ and $\tilde{H}_*(L_\alpha)$ in $\dim \leq n - 1$. Note that $H_n(L) \otimes \mathbb{Z}_{(\mathcal{A})} = H_n(L) \otimes \mathbb{Z}[1/q] = (H_n(L)/\text{Tor}_q H_n(L)) \otimes \mathbb{Z}[1/q]$ and since $H_n(L)/\text{Tor}_q H_n(L)$ is q -divisible we have that the $\mathcal{A}$ -localization of $H_n(L)$ is an epimorphism. Then by the Whitehead theorem α induces an isomorphism of $\pi_i(L)$ and $\pi_i(L_\alpha)$, $i \leq n - 1$ and an epimorphism of $\pi_n(L)$ and $\pi_n(L_\alpha)$. Therefore $\pi_i(L)$, $i \leq n - 1$ is $\mathcal{A}$ -local (that is, q -divisible and q -torsion free) and the $\mathcal{A}$ -localization of $\pi_n(L)$ is an epimorphism. The last property implies that $\pi_n(L)/\text{Tor}_q \pi_n(L)$ is q -divisible.

Let $q \in \mathcal{P}$, $q \neq p$ and $q \notin \mathcal{F}$.

Let $\beta : L \longrightarrow L_\beta$ be a q -completion of L . Then β induces an isomorphism of $\tilde{H}_*(L; \mathbb{Z}_q)$ and $\tilde{H}_*(L_\beta; \mathbb{Z}_q)$ and since $H_n(L; \mathbb{Z}_q) = 0$ we get that $H_n(L_\beta; \mathbb{Z}_q) = 0$ and therefore $H_n(L_\beta)$ is q -divisible. Now since $\pi_i(L)$ is q -divisible and q -torsion free we have that $\text{Hom}(\mathbb{Z}_{q^\infty}, \pi_i(L)) = 0$, $i \leq n - 1$ and by Proposition 2.1 $\text{Ext}(\mathbb{Z}_{q^\infty}, \pi_i(L)) = 0$, $i \leq n - 1$. Then the exact sequence

$$0 \longrightarrow \text{Ext}(\mathbb{Z}_{q^\infty}, \pi_i(L)) \longrightarrow \pi_i(L_\beta) \longrightarrow \text{Hom}(\mathbb{Z}_{q^\infty}, \pi_{i-1}(L)) \longrightarrow 0$$

implies that L_β is $(n-1)$ -connected and $\text{Ext}(\mathbb{Z}_{q^\infty}, \pi_n(L)) = \pi_n(L_\beta)$. Thus $\pi_n(L_\beta) = H_n(L_\beta)$ and hence $\text{Ext}(\mathbb{Z}_{q^\infty}, \pi_n(L))$ is q -divisible. Then by Proposition 2.1 $\pi_n(L)$ is q -divisible and (iii) is proved.

(iv). The proof is similar to the proof of (iii). □

Let X be a compactum and let n be a positive integer. The Bockstein basis of abelian groups is the following collection of groups $\sigma = \{\mathbb{Q}, \mathbb{Z}_p, \mathbb{Z}_{p^\infty}, \mathbb{Z}_{(p)} : p \in \mathcal{P}\}$. Define the Bockstein basis $\sigma(X, n)$ of X in dimensions $\leq n$ as $\sigma(X, n) = \{E \in \sigma : \dim_E X \leq n\}$. Following [6] denote:

$$\mathcal{T}(X, n) = \{p \in \mathcal{P} : \mathbb{Z}_p \text{ or } \mathbb{Z}_{p^\infty} \in \sigma(X, n)\} ;$$

$$\mathcal{D}(X, n) = \emptyset \text{ if } \sigma(X, n) \text{ contains only torsion groups;}$$

$$\mathcal{D}(X, n) = \mathcal{P} \text{ if } \mathbb{Q} \in \sigma(X, n) \text{ but none of } \mathbb{Z}_{(p)}, p \in \mathcal{P} \text{ belongs to } \sigma(X, n) \text{ and}$$

$$\mathcal{D}(X, n) = \mathcal{P} \setminus \{p \in \mathcal{P} : \mathbb{Z}_{(p)} \in \sigma(X, n)\} \text{ otherwise ;}$$

$$\mathcal{F}(X, n) = \mathcal{D}(X, n) \setminus \mathcal{T}(X, n).$$

Note that for every group G such that $\dim_G X \leq n$, G is $\mathcal{F}(X, n)$ -torsion free.

Proposition 2.3 *Let X be a compactum such that $\mathcal{D}(X, n) \neq \emptyset$. Then $\dim_H X \leq n$ for every group H such that H is $\mathcal{D}(X, n)$ -divisible and $\mathcal{F}(X, n)$ -torsion free.*

Proof. Let $G = \oplus\{E : E \in \sigma(X, n)\}$. Then $\dim_G X \leq n$. One can easily verify that in the notations of Proposition 2.4 of [6], $\mathcal{D}(G) = \mathcal{D}(X, n)$ and $\mathcal{F}(G) = \mathcal{F}(X, n)$. Then the result follows from Proposition 2.4 of [6]. □

In the proof of Theorem 1.1 we will also use the following facts.

Proposition 2.4 ([7]) *Let K be a simply connected CW-complex such that K has only finitely many non-trivial homotopy groups. Let X be a compactum such that $\dim_{\pi_i(K)} X \leq i$ for $i > 1$. Then $e - \dim X \leq K$.*

Let K' be a simplicial complex. We say that maps $h : K \longrightarrow K'$, $g : L \longrightarrow L'$, $\alpha : L \longrightarrow K$ and $\alpha' : L' \longrightarrow K'$ combinatorially commute if for every simplex Δ of K' we have that $(\alpha' \circ g)((h \circ \alpha)^{-1}(\Delta)) \subset \Delta$. Recall that a map $h' : K \longrightarrow L'$ is a combinatorial lifting of h to L' if for every simplex Δ of K' we have that $(\alpha' \circ h')(h^{-1}(\Delta)) \subset \Delta$.

For a simplicial complex K and $a \in K$, $st(a)$ denotes the union of all the simplices of K containing a .

Proposition 2.5 ([7])

(i) Let a compactum X be represented as the inverse limit $X = \varprojlim K_i$ of finite simplicial complexes K_i with bonding maps $h_j^i : K_j \rightarrow K_i$. Fix i and let $\omega : EW(K_i, k) \rightarrow K_i$ be a resolution of K_i which is suitable for X . Then there is a sufficiently large j such that h_j^i admits a combinatorial lifting to $EW(K_i, k)$.

(ii) Let $h : K \rightarrow K'$, $h' : K \rightarrow L'$ and $\alpha' : L' \rightarrow K'$ be maps of a simplicial complex K' and CW-complexes K and L' such that h and α' are combinatorial and h' is a combinatorial lifting of h . Then there is a cellular approximation of h' which is also a combinatorial lifting of h .

(iii) Let K and K' be simplicial complexes, let maps $h : K \rightarrow K'$, $g : L \rightarrow L'$, $\alpha : L \rightarrow K$ and $\alpha' : L' \rightarrow K'$ combinatorially commute and let h be combinatorial. Then

$g(\alpha^{-1}(st(x))) \subset \alpha'^{-1}(st(h(x)))$ and $h(st((\alpha(z)))) \subset st((\alpha' \circ g)(z))$ for every $x \in K$ and $z \in L$;

3 Proof of Theorem 1.1

Denote $\mathcal{D} = \mathcal{D}(X, n)$ and $\mathcal{F} = \mathcal{F}(X, n)$. Represent X as the inverse limit $X = \varprojlim (K_i, h_i)$ of finite simplicial complexes K_i with combinatorial bonding maps $h_{i+1} : K_{i+1} \rightarrow K_i$ and the projections $p_i : X \rightarrow K_i$ such that for every simplex Δ of K_i , $\text{diam}(p_i^{-1}(\Delta)) \leq 1/i$. Following A. Dranishnikov [3, 4] we construct by induction finite CW-complexes L_i and maps $g_{i+1} : L_{i+1} \rightarrow L_i$, $\alpha_i : L_i \rightarrow K_i$ such that

(a) L_i is $(n+1)$ -dimensional and obtained from $K_i^{[n+1]}$ by replacing some $(n+1)$ -simplexes by $(n+1)$ -cells attached to the boundary of the replaced simplexes by a map of degree $\in S(\mathcal{F})$. Then α_i is a projection of L_i taking the new cells to the original ones such that α_i is 1-to-1 over $K_i^{[n]}$. We define a simplicial structure on L_i for which α_i is a combinatorial map and refer to this simplicial structure while constructing resolutions of L_i . Note that for $\mathcal{F} = \emptyset$ we don't replace simplexes of $K_i^{[n+1]}$ at all;

(b) the maps h_{i+1} , g_{i+1} , α_{i+1} and α_i combinatorially commute. Recall that this means that for every simplex Δ of K_i , $(\alpha_i \circ g_{i+1})((h_{i+1} \circ \alpha_{i+1})^{-1}(\Delta)) \subset \Delta$.

We will construct L_i in such a way that $Z = \varprojlim (L_i, g_i)$ will admit a map $r : Z \rightarrow X$ such that Z and r satisfy the conclusions of the theorem.

Let $E \in \sigma$ be such that $\dim_E X \leq k$, $2 \leq k \leq n$ and let $f : N \rightarrow K(E, k)$ be a cellular map from a subcomplex N of L_i , $L_i^{[k]} \subset N$. Let $\omega_L : EW(L_i, k) \rightarrow L_i$ be the standard resolution of L_i for f . We are going to construct from $\omega_L : EW(L_i, k) \rightarrow L_i$ a resolution $\omega : EW(K_i, k) \rightarrow K_i$ of K_i suitable for X . If $\dim K_i \leq k$ set $\omega = \alpha_i \circ \omega_L : EW(K_i, k) = EW(L_i, k) \rightarrow K_i$.

If $\dim K_i > k$ set $\omega_k = \alpha_i \circ \omega_L : EW_k(K_i, k) = EW(L_i, k) \longrightarrow K_i$ and we will construct by induction resolutions $\omega_j : EW_j(K_i, k) \longrightarrow K_i$, $k+1 \leq j \leq \dim K_i$ such that $EW_j(K_i, k)$ is a subcomplex of $EW_{j+1}(K_i, k)$ and ω_{j+1} extends ω_j for every $k \leq j < \dim K_i$.

Assume that $\omega_j : EW_j(K_i, k) \longrightarrow K_i$, $k \leq j < \dim K_i$ is constructed. For every simplex Δ of K_i of $\dim = j+1$ consider the subcomplex $\omega_j^{-1}(\Delta)$ of $EW_j(K_i, k)$. Enlarge $\omega_j^{-1}(\Delta)$ by attaching cells of $\dim = n+1$ in order to kill the elements of $\text{Tor}_{\mathcal{F}} \pi_n(\omega_j^{-1}(\Delta))$ and attaching cells of $\dim > n+1$ in order to get a subcomplex with trivial homotopy groups in $\dim > n$. Let $EW_{j+1}(K_i, k)$ be $EW_j(K_i, k)$ with all the cells attached for all $(j+1)$ -dimensional simplexes Δ of K_i and let $\omega_{j+1} : EW_{j+1}(K_i, k) \longrightarrow K_i$ be an extension of ω_j sending the interior points of the attached cells to the interior of the corresponding Δ .

Finally denote $EW(K_i, k) = EW_j(K_i, k)$ and $\omega = \omega_j : EW_j(K_i, k) \longrightarrow K_i$ for $j = \dim K_i$. Note that since we attach cells only of $\dim > n$, the n -skeleton of $EW(K_i, k)$ coincides with the n -skeleton of $EW(L_i, k)$.

Let us show that $EW(K_i, k)$ is suitable for X . Fix a simplex Δ of K_i . Since $\omega^{-1}(\Delta)$ is contractible if $\dim \Delta \leq k$ assume that $\dim \Delta > k$. Denote $T = \alpha_i^{-1}(\Delta)$. Note that it follows from the construction that T is $(n-1)$ -connected, $H_n(T)$ is $\mathcal{F}$ -torsion, $\omega^{-1}(\Delta)$ is $(k-1)$ -connected, $\pi_n(\omega^{-1}(\Delta)) = \pi_n(\omega_L^{-1}(T))/\text{Tor}_{\mathcal{F}} \pi_n(\omega_L^{-1}(T))$, $\pi_j(\omega^{-1}(\Delta)) = 0$ for $j \geq n+1$ and $\pi_j(\omega^{-1}(\Delta)) = \pi_j(\omega_L^{-1}(T))$ for $j \leq n-1$.

Consider the following cases.

Case 1. $E = \mathbb{Z}_p$. By (i) of Proposition 2.2, $\pi_j(\omega_L^{-1}(T))$, $j \leq n-1$ and $\pi_n(\omega_L^{-1}(T))/\text{Tor}_{\mathcal{F}} \pi_n(\omega_L^{-1}(T))$ are p -torsion. Then $\pi_j(\omega^{-1}(\Delta))$ is p -torsion for $j \leq n$. Therefore $\dim_{\pi_j(\omega^{-1}(\Delta))} X \leq \dim_{\mathbb{Z}_p} X \leq k$ for $j \geq k$ and hence by Proposition 2.4 e - $\dim X \leq \omega^{-1}(\Delta)$.

Case 2. $E = \mathbb{Z}_{p^\infty}$. Then by (ii) of Proposition 2.2, $\pi_j(\omega_L^{-1}(T))$, $j \leq n-1$ and $\pi_n(\omega_L^{-1}(T))/\text{Tor}_{\mathcal{F}} \pi_n(\omega_L^{-1}(T))$ are p -torsion and $\pi_k(\omega_L^{-1}(T))$ is p -divisible. Then $\pi_j(\omega^{-1}(\Delta))$ is p -torsion for $j \leq n$ and $\pi_k(\omega^{-1}(\Delta))$ is p -divisible. Therefore by the Bockstein theorem and inequalities $\dim_{\pi_k(\omega^{-1}(\Delta))} X \leq \dim_{\mathbb{Z}_{p^\infty}} X \leq k$ and $\dim_{\pi_j(\omega^{-1}(\Delta))} X \leq \dim_{\mathbb{Z}_{p^\infty}} X + 1 \leq k+1$ for $j \geq k+1$. Hence by Proposition 2.4 e - $\dim X \leq \omega^{-1}(\Delta)$.

Case 3. $E = \mathbb{Z}_{(p)}$. Then by (iii) of Proposition 2.2, $\pi_j(\omega_L^{-1}(T))$, $j \leq n-1$ is p -local and $\pi_n(\omega_L^{-1}(T))/\text{Tor}_{\mathcal{F}} \pi_n(\omega_L^{-1}(T))$ is q -divisible for every $q \in \mathcal{P}$, $q \neq p$. Then $\pi_j(\omega^{-1}(\Delta))$, $j \leq n-1$ is p -local and $\pi_n(\omega^{-1}(\Delta))$ is $\mathcal{D}$ -divisible and $\mathcal{F}$ -torsion free. Therefore $\dim_{\pi_j(\omega^{-1}(\Delta))} X \leq k$ for $j \leq n-1$ and by Proposition 2.3, $\dim_{\pi_n(\omega^{-1}(\Delta))} X \leq n$. Hence by Proposition 2.4 e - $\dim X \leq \omega^{-1}(\Delta)$.

Case 4. $E = \mathbb{Q}$. This case is similar to the previous one.

Thus we have shown that $EW(K_i, k)$ is suitable for X . Now replacing K_{i+1} by K_j with a sufficiently large j we may assume by (i) of Proposition 2.5 that there is

a combinatorial lifting of h_{i+1} to $h'_{i+1} : K_{i+1} \longrightarrow EW(K_i, k)$. By (ii) of Proposition 2.5 we replace h'_{i+1} by its cellular approximation preserving the property of h'_{i+1} of being a combinatorial lifting of h_{i+1} .

Let Δ be a simplex of K_i and let $\tau : (\alpha_i \circ \omega_L)^{-1}(\Delta \longrightarrow \omega^{-1}(\Delta))$ be the inclusion. Note that from the construction it follows that for the induced homomorphism $\tau_* : \pi_n((\alpha_i \circ \omega_L)^{-1}(\Delta)) \longrightarrow \pi_n(\omega^{-1}(\Delta))$, $\ker \tau_*$ is $\mathcal{F}$ -torsion. Using this fact and the reasoning described in detail in the proof of Theorem 1.2 of [6] one can construct from $K_{i+1}^{[n+1]}$ a CW-complex L_{i+1} by replacing some $(n+1)$ -simplexes of $K_{i+1}^{[n+1]}$ by $(n+1)$ -cells attached to the boundary of the replaced simplexes by a map of degree $\in S(\mathcal{F})$ such that h'_{i+1} restricted to $K_{i+1}^{[n]}$ extends to a map $g'_{i+1} : L_{i+1} \longrightarrow EW(L_i, n)$ such that g'_{i+1} , α_{i+1} , h_{i+1} and $\alpha_i \circ \omega_L$ combinatorially commute where α_{i+1} is a projection of L_{i+1} into K_{i+1} taking the new cells to the original ones such that α_{i+1} is 1-to-1 over $K_{i+1}^{[n]}$.

Now define $g_{i+1} = \omega_L \circ g'_{i+1} : L_{i+1} \longrightarrow L_i$ and finally define a simplicial structure on L_{i+1} for which α_{i+1} is a combinatorial map. It is easy to check that the properties (a) and (b) are satisfied. Since the triangulation of L_{i+1} can be replaced by any of its barycentric subdivisions we may also assume that

(c) $\text{diam} g_{i+1}^j(\Delta) \leq 1/i$ for every simplex Δ in L_{i+1} and $j \leq i$
 where $g_i^j = g_{j+1} \circ g_{j+2} \circ \dots \circ g_i : L_i \longrightarrow L_j$.

Denote $Z = \varprojlim (L_i, g_i)$ and let $r_i : Z \longrightarrow L_i$ be the projections.

Clearly $\dim Z \leq n+1$. For constructing L_{i+1} we used an arbitrary map $f : N \longrightarrow K(E, k)$ such that $E \in \sigma$, $\dim_E X \leq k$, $2 \leq k \leq n$ and N is a subcomplex of L_i containing $L_i^{[k]}$. By a standard reasoning described in detail in the proof of Theorem 1.6 of [7] one can show that choosing E and f in an appropriate way for each i we can achieve that $\dim_E Z \leq k$ for every integer $2 \leq k \leq n$ and every $E \in \sigma$ such that $\dim_{\mathbb{Z}_p} X \leq k$. Then by the Bockstein theorem $\dim_G Z \leq k$ for every group G such that $\dim_G X \leq k$, $2 \leq k \leq n$.

By (iii) of Proposition 2.5, the properties (a) and (b) imply that for every $x \in X$ and $z \in Z$ the following holds:

- (d1) $g_{i+1}(\alpha_{i+1}^{-1}(st(p_{i+1}(x)))) \subset \alpha_i^{-1}(st(p_i(x)))$ and
- (d2) $h_{i+1}(st((\alpha_{i+1} \circ r_{i+1})(z))) \subset st((\alpha_i \circ r_i)(z))$.

Define a map $r : Z \longrightarrow X$ by $r(z) = \cap \{p_i^{-1}(st((\alpha_i \circ r_i)(z))) : i = 1, 2, \dots\}$. Then (d2) implies that r is indeed well-defined and continuous.

The properties (d1) and (d2) also imply that for every $x \in X$

$$r^{-1}(x) = \varprojlim (\alpha_i^{-1}(st(p_i(x))), g_i|_{\alpha_i^{-1}(st(p_i(x)))})$$

where the map $g_i|_{\dots}$ is considered as a map to $\alpha_{i-1}^{-1}(st(p_{i-1}(x)))$.

Since $r^{-1}(x)$ is not empty for every $x \in X$, r is a map onto. Fix $x \in X$ and let us show that $r^{-1}(x)$ satisfies the conclusions of the theorem. Since $T = \alpha_i^{-1}(st(p_i(x)))$ is $(n-1)$ -connected we obtain that $r^{-1}(x)$ is approximately $(n-1)$ -connected as the inverse limit of $(n-1)$ -connected finite simplicial complexes.

Let a group G be such that $\dim_G X \leq n$. Note that $H_n(T)$ is $\mathcal{F}$ -torsion and G is $\mathcal{F}$ -torsion free. Then by the universal-coefficient theorem $H^n(T; G) = \text{Hom}(H_n(T), G) = 0$. Thus $\tilde{H}^k(r^{-1}(x); G) = 0$ for $k \leq n$ and since $\dim_G Z \leq n$, $\tilde{H}^k(r^{-1}(x); G) = 0$ for $k \geq n+1$. Hence r is G -acyclic and this completes the proof. $\square$

References

- [1] Bousfield, A. K.; Kan, D. M. Homotopy limits, completions and localizations. Lecture Notes in Mathematics, Vol. 304, Springer-Verlag, Berlin-New York, 1972.
- [2] M. Cencelj and A.N. Dranishnikov, Extension of maps to nilpotent spaces. II, Topology Appl. 124(2002), no. 1, 77-83.
- [3] Dranishnikov, A. N. Rational homology manifolds and rational resolutions. Special issue in memory of B. J. Ball. Topology Appl. 94 (1999), no. 1-3, 75-86.
- [4] A. Dranishnikov, Cohomological dimension theory of compact metric spaces, Topology Atlas Invited Contributions, <http://at.yorku.ca/topology/taic.htm>
- [5] A. Koyama and K. Yokoi, Cohomological dimension and acyclic resolutions. In memory of T. Benny Rushing. Topology Appl. 120 (2002), no. 1-2, 175-204.
- [6] M. Levin, Acyclic resolutions for arbitrary groups, Israel J. Math. 135(2003), 193-203.
- [7] M. Levin, Cell-like resolutions preserving cohomological dimensions, Algebr. Geom. Topol. 3(2003), 1277-1289.
- [8] M. Levin, Universal acyclic resolutions for finitely generated coefficient groups, Topology Appl. 135(2004), 101-109.

Department of Mathematics
Ben Gurion University of the Negev
P.O.B. 653
Be'er Sheva 84105, ISRAEL
e-mail: mlevine@math.bgu.ac.il

